

First aid fact sheet

Choking adult or child (over 1 year)



If the patient becomes blue, limp or unconscious, follow DRSABCD and call **Triple Zero (000)** for an ambulance.

Signs and symptoms

- clutching the throat
- coughing, wheezing, gagging
- difficulty in breathing, speaking or swallowing
- making a whistling or 'crowing' noise, or no sound at all
- blue lips, face, earlobes, fingernails
- loss of consciousness



5 back blows



5 chest thrusts

What to do

- 1 Encourage the patient to relax. Ask the patient to cough to remove the object.
- 2 If coughing does not remove the blockage, call **Triple Zero (000)** for an ambulance.
- 3 Bend the patient well forward and give up to 5 sharp blows on the back between the shoulder blades with the heel of one hand.
- 4 Check if the blockage has been removed after each blow.
- 5 If the blockage has not cleared after 5 back blows, give up to 5 chest thrusts.
- 6 Give chest thrusts by placing one hand in the middle of the patient's back for support and the heel of the other on the lower half of the sternum. Thrusts should be slower and sharper than CPR compressions.
- 7 Check if the blockage has been removed after each thrust.
- 8 If the blockage has not cleared after 5 thrusts, continue alternating 5 back blows with 5 chest thrusts until medical aid arrives.
- 9 If the patient becomes blue, limp or unconscious, follow DRSABCD and call **Triple Zero (000)** for an ambulance.

In a medical emergency call Triple Zero (000)

DRSABCD Danger ► Response ► Send for help ► Airway ► Breathing ► CPR ► Defibrillation

You could save a life with first aid training • www.stjohn.org.au • **1300 360 455**

This information is not a substitute for first aid training. Formal instruction in resuscitation is essential. St John Ambulance Australia is not liable for any damages or incidents that may occur in the use of this information by other parties or individuals. This is not for commercial distribution. © St John Ambulance Australia 2022.