



## First aid fact sheet

# Febrile convulsion



- A rapid rise in body temperature can cause convulsions. This occurs most often in children aged 6 months to 5 years.
- A convulsion can occur with a temperature change from the normal 37°C to as little as 38.5°C.

### Signs and symptoms

Symptoms can include:

- fever
- muscle stiffening
- twitching or jerking of face or limbs
- eyes rolling upwards
- blue lips, face, earlobes, fingernails
- loss of consciousness

### What to do

#### During the convulsion

- 1 Place the child on their side for safety.
- 2 DO NOT restrain the child.
- 3 DO NOT cool the child by sponging or bathing, but do remove excess clothing.

#### After the convulsion

- 4 Follow DRSABCD.
- 5 Remove excess clothing or wrappings.
- 6 Seek medical aid.

**In a medical emergency call Triple Zero (000)**

**DRSABCD** Danger ► Response ► Send for help ► Airway ► Breathing ► CPR ► Defibrillation

You could save a life with first aid training • [www.stjohn.org.au](http://www.stjohn.org.au) • **1300 360 455**

This information is not a substitute for first aid training. Formal instruction in resuscitation is essential. St John Ambulance Australia is not liable for any damages or incidents that may occur in the use of this information by other parties or individuals. This is not for commercial distribution. © St John Ambulance Australia 2022.